

Diemer F, Sutor V. Praxis der Medizinischen Trainingstherapie II. Stuttgart: Thieme; 2010

Literaturliste zu Kapitel 4:

Alpert SW, Pink MM, Jobe FW, McMahon PJ, Mathiyakom W. Electromyographic analysis of deltoid and rotator cuff function under varying loads and speeds. J Shoulder Elbow Surg. 2000 Jan-Feb;9(1):47.

Barnett C, Kippers V, Turner P. Effects of variations of the bench press exercise on the EMG activity of five shoulder muscles. J Strength Condition Res. 1995 9: 222.

Barnett C, Kippers V, Turner P. Effects of Variations of the Bench Press Exercise on the EMG Activity of Five Shoulder Muscles. Journal of Strength & Conditioning Research. 1995 9(4):222.

Bisset LM, Coppieters MW, Vicenco B. Sensimotor deficits remain despite resolution of symptoms using conservative treatment in patients with tennis elbow: a randomized controlled trial. Archives of Physical Medicine and Rehabilitation. 2009; 90: 1.

Boeckh-Behrens UW. Supertrainer Schultern, Arme, Brust: Die effektivsten Übungen. Die besten Programme. Wissenschaftlich getestet. rororo; 3.Auflage. 2005

Borstad JD, Ludewig PM. Comparison of three stretches for the pectoralis minor muscle. J Shoulder Elbow Surg. 2006 May-Jun;15(3):324.

Borstad JD. Measurement of pectoralis minor muscle length: validation and clinical application. J Orthop Sports Phys Ther. 2008 Apr;38(4):169.

Borstad JD. Measurement of pectoralis minor muscle length: validation and clinical application. J Orthop Sports Phys Ther. 2008 Apr;38(4):169.

Bradley JP, Tibone JE. Electromyographic analysis of muscle action about the shoulder. Clin Sports Med. 1991 Oct;10(4):789.

Cagnie B, Danneels L, Cools A et al. The influence of breathing type, expiration and cervical posture on the performance of the cranio-cervical flexion test in healthy subjects. Manual Therapy. 2008; 13: 232.

Clisby EF, Bitter NL, Sandow MJ, Jones MA, Magarey ME, Jaberzadeh S. Relative contributions of the infraspinatus and deltoid during external rotation in patients with symptomatic subacromial impingement. J Shoulder Elbow Surg. 2008 Jan-Feb;17(1 Suppl):87S.

Cogley RM, Archambault TA, Fibeger JF, Koverman MM, Youdas JW, Hollman JH. Comparison of muscle activation using various hand positions during the push-up exercise. J Strength Cond Res. 2005 Aug;19(3):628.

Cools AM, Dewitte V, Lanszweert F, Notebaert D, Roets A, Soetens B, Cagnie B, Witvrouw EE. Rehabilitation of scapular muscle balance: which exercises to prescribe? Am J Sports Med. 2007 Oct;35(10):1744.

De Souza Castelo Oliviera A, Goncalves M. Lumbar muscles recruitment during resistance exercise for upper limbs. *Journal of Electromyography and Kinesiology*. 2008; published online.

Decker MJ, Tokish JM, Ellis HB, Torry MR, Hawkins RJ. Subscapularis muscle activity during selected rehabilitation exercises. *Am J Sports Med*. 2003 Jan-Feb;31(1):126.

Demaille-Wlodyka S, Chiquet C, Lavaste JF et al. Cervical range of motion and cephalic kinesthesia. *Spine*. 2007 ; 32 : E254.

Ebaugh DD, McClure PW, Karduna AR. Scapulothoracic and glenohumeral kinematics following an external rotation fatigue protocol. *J Orthop Sports Phys Ther*. 2006 Aug;36(8):557.

Ekstrom RA, Donatelli RA, Soderberg GL. Surface electromyographic analysis of exercises for the trapezius and serratus anterior muscles. *J Orthop Sports Phys Ther*. 2003 May;33(5):247.

Ellenbecker TS. *Shoulder Rehabilitation*. Thieme. New York. 2006.

Falla D, O`Leary S, Fagan A et al. Recruitment of the deep cervical flexor muscles during a postural-correction exercise performed in sitting. *Manual Therapy*. 2007; 12: 139.

Falla D. Muskeln effektiv kontrollieren. *Physiopraxis*. 2006; 3: 18.

Gokeler A, van Paridon-Edauw GH, DeClercq S, Matthijs O, Dijkstra PU. Quantitative analysis of traction in the glenohumeral joint. In vivo radiographic measurements. *Man Ther*. 2003 May;8(2):97.

Graichen H, Bonel H, Stammberger T, Haubner M, Rohrer H, Englmeier KH, Reiser M, Eckstein F. Three-dimensional analysis of the width of the subacromial space in healthy subjects and patients with impingement syndrome. *AJR Am J Roentgenol*. 1999 Apr;172(4):1081.

Graichen H, Bonel H, Stammberger T, Heuck A, Englmeier KH, Reiser M, Eckstein F. An MR-based technique for determination of the subacromial space width in subjects with and without shoulder muscle activity. *Z Orthop Ihre Grenzgeb*. 1999 Jan-Feb;137(1):2.

Graichen H, Hinterwimmer S, von Eisenhart-Rothe R, Vogl T, Englmeier KH, Eckstein F. Effect of abducting and adducting muscle activity on glenohumeral translation, scapular kinematics and subacromial space width in vivo. *J Biomech*. 2005 Apr;38(4):755.

Hill R, Jensen P, Baardsen T et al. Head repositioning accuracy to neutral: a comparative study of error calculation. *Manual Therapy*. 2009; 14: 110.

Hintermeister RA, Lange GW, Schultheis JM, Bey MJ, Hawkins RJ. Electromyographic activity and applied load during shoulder rehabilitation exercises using elastic resistance. *Am J Sports Med*. 1998 Mar-Apr;26(2):210.

Hinterwimmer S, Von Eisenhart-Rothe R, Siebert M, Putz R, Eckstein F, Vogl T, Graichen H. Influence of adducting and abducting muscle forces on the subacromial space width. *Med Sci Sports Exerc*. 2003 Dec;35(12):2055.

Jull GA, O`Leary SP, Falla DL. Clinical assessment of the deep cervical flexor muscles: the craiocervical flexion test. *Journal of Manipulative and Physiological Therapeutics*. 2008; 31: 525.

Juul-Kristensen B, Lund H, Hansen K et al. Poorer elbow proprioception in patients with lateral epicondylitis than in healthy controls: a cross-sectional study. 2008; *Journal of Elbow and Shoulder Surgery*. 2008; 17: 72.

Kelly BT, Kadrmas WR, Speer KP. The manual muscle examination for rotator cuff strength. An electromyographic investigation. *Am J Sports Med*. 1996 Sep-Oct;24(5):581.

Kibler WB, Uhl TL, Maddux JW, Brooks PV, Zeller B, McMullen J. Qualitative clinical evaluation of scapular dysfunction: a reliability study. *J Shoulder Elbow Surg*. 2002 Nov-Dec;11(6):550.

Kibler WB. Scapular involvement in impingement: signs and symptoms. *Instr Course Lect*. 2006;55:35.

Koslow PA, Prosser LA, Strony GA, Suchecki SL, Mattingly GE. Specificity of the lateral scapular slide test in asymptomatic competitive athletes. *J Orthop Sports Phys Ther*. 2003 Jun;33(6):331.

Kramer M et al. Reliability of a new virtual reality test to measure cervicocephalic kinesthesia. *Journal of Electromyography and Kinesiology*. 2008. doi: 10.1016/j.jelekin. 2008.05.005.

Kristjansson E, Hardardottir L, Asmundardottir M et al. a new clinical test for cervicocephalic kinesthetic sensibility: the fly. *Archives of Physical Medicine and Rehabilitation*. 2004; 85: 490.

Lagally KM, McCaw ST, Young GT, Medema HC, Thomas DQ. Ratings of perceived exertion and muscle activity during the bench press exercise in recreational and novice lifters. *J. Strength Cond. Res*. 2004. 18:359.

Lee HY, Teng CC, Chai HM et al. Test-retest reliability of cervicocephalic kinesthetic sensibility in three cardinal planes. *Manual Therapy*. 2006; 11: 61.

Lehman GJ. The Influence of Grip Width and Forearm Pronation/Supination on Upper-Body Myoelectric Activity During the Flat Bench Press. *Journal of Strength & Conditioning Research*. 2005 19(3):587.

Lewis JS, Valentine RE. The pectoralis minor length test: a study of the intra-rater reliability and diagnostic accuracy in subjects with and without shoulder symptoms. *BMC Musculoskelet Disord*. 2007 Jul 9;8:64.

Magarey ME, Jones MA. Dynamic evaluation and early management of altered motor control around the shoulder complex. *Man Ther*. 2003 Nov;8(4):195.

Magarey ME, Jones MA. Specific evaluation of the function of force couples relevant for stabilization of the glenohumeral joint. *Man Ther*. 2003 Nov;8(4):247.

McClure P, Balaicuis J, Heiland D, Broersma ME, Thorndike CK, Wood A. A randomized controlled comparison of stretching procedures for posterior shoulder tightness. *J Orthop Sports Phys Ther*. 2007 Mar;37(3):108.

Myers JB, Pasquale MR, Laudner KG, Sell TC, Bradley JP, Lephart SM. On-the-Field Resistance-Tubing Exercises for Throwers: An Electromyographic Analysis. *J Athl Train*. 2005 Mar;40(1):15.

Nijs J, Roussel N, Vermeulen K, Souvereyns G. Scapular positioning in patients with shoulder pain: a study examining the reliability and clinical importance of 3 clinical tests. *Arch Phys Med Rehabil.* 2005;86(7):1349.

Odom CJ, Taylor AB, Hurd CE, Denegar CR. Measurement of scapular asymmetry and assessment of shoulder dysfunction using the Lateral Scapular Slide Test: a reliability and validity study. *Phys Ther.* 2001 Feb;81(2):799.

Pinsault N, Vuillerme N, Pavan P. Cervicocephalic reolocation test to the neutral head position : assessment in bilateral labyrinthine-defective and chronic, nontraumatic neck pain patients. *Archives of Physical Medicine and Rehabilitation.* 2008; 89: 2375.

Rabin A, Irrgang JJ, Fitzgerald GK, Eubanks A. The intertester reliability of the Scapular Assistance Test. *J Orthop Sports Phys Ther.* 2006 Sep;36(9):653.

Reddy AS, Mohr KJ, Pink MM, Jobe FW. Electromyographic analysis of the deltoid and rotator cuff muscles in persons with subacromial impingement. *J Shoulder Elbow Surg.* 2000 Nov-Dec;9(6):519.

Reinold MM, Macrina LC, Wilk KE, Fleisig GS, Dun S, Barrentine SW, Ellerbusch MT, Andrews JR. Electromyographic analysis of the supraspinatus and deltoid muscles during 3 common rehabilitation exercises. *J Athl Train.* 2007 Oct-Dec;42(4):464.

Revel M, Andre-Deshays CA, Minguet M. Cervicocephalic kinesthetic sensibility in patients with cervical pain. *Archives of Physical Medicine and Rehabilitation.* 1991; 72: 288.

Revel M, Minguet M, Gergory P et al. Changes in Cervicocephalic kinesthesia after a proprioceptive rehabilitation program in patients with neck pain. *Archives of Physical Medicine and Rehabilitation.* 1994; 75: 895.

Rix GD, Bagust J. Cervicocephalic kinesthetic sensibility in patients with chronic, nontraumatic cervical spine pain. *Archives of Physical Medicine and Rehabilitation.* 2001; 82: 911.

Sahara W, Sugamoto K, Murai M, Tanaka H, Yoshikawa H. 3D kinematic analysis of the acromioclavicular joint during arm abduction using vertically open MRI. *J Orthop Res.* 2006 Sep;24(9):1823.

Sahrmann SA. Movement impairment syndromes of the shoulder girdle. In: *Diagnosis and treatment of movement impairment syndromes.* St Louis: Mosby; 2002. p. 193.

Santana JC, Vera-Garcia FJ, McGill SM. A kinetic and electromyographic comparison of the standing cable press and bench press. *J Strength Cond Res.* 2007 Nov;21(4):1271.

Schwellnus MP. The repeatability of clinical and laboratory tests to measure scapular position and movement during arm abduction. *Intl Sports Med J.* 2003;4:1.

Shadmehr A, Bagheri H, Ansari NN, Sarafraz H. The reliability measurements of lateral scapular slide test at three different degrees of shoulder joint abduction. *Br J Sports Med.* 2010 Mar;44(4):289.

Signorile JF, Zink AJ, Szwed SP. A comparative electromyographical investigation of muscle utilization patterns using various hand positions during the lat pull-down. *J Strength Cond Res.* 2002 Nov;16(4):539.

Treleaven J, Jull G, LowChoy N. The relationship of cervical joint position error to balance and eye movement disturbances in persistent whiplash. *Manual Therapy.* 2006; 11: 99.

Treleaven J. Sensimotor disturbances in neck disorders affecting postural stability, head and eye movement control. *Manual Therapy.* 2008a; 13: 2.

Treleaven J. Sensimotor disturbances in neck disorders affecting postural stability, head and eye movement control – part 2: case studies. *Manual Therapy.* 2008b; 13: 266.

Uhl TL, Carver TJ, Mattacola CG, Mair SD, Nitz AJ. Shoulder musculature activation during upper extremity weight-bearing exercise. *J Orthop Sports Phys Ther.* 2003 Mar;33(3):109.